

D-equivalence and L-equivalence of K3 surfaces

Evgeny Shinder (Sheffield)

Derived category and birational geometry, Osaka
21.02.2017

Introduction: Geometric meaning of D-equivalence

Grothendieck ring of varieties and L-equivalence

Quadrics, quadric fibrations and K3 surfaces

D-equivalence

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

- ▶ Mukai: if A abelian variety, then $\mathcal{D}^b(A) \simeq \mathcal{D}^b(\widehat{A})$

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

- ▶ Mukai: if A abelian variety, then $\mathcal{D}^b(A) \simeq \mathcal{D}^b(\widehat{A})$
- ▶ Abelian varieties [complete classification of D-equivalence by Orlov]

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

- ▶ Mukai: if A abelian variety, then $\mathcal{D}^b(A) \simeq \mathcal{D}^b(\widehat{A})$
- ▶ Abelian varieties [complete classification of D-equivalence by Orlov]
- ▶ K3 surfaces [complete classification of D-equivalence by Mukai]

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

- ▶ Mukai: if A abelian variety, then $\mathcal{D}^b(A) \simeq \mathcal{D}^b(\widehat{A})$
- ▶ Abelian varieties [complete classification of D-equivalence by Orlov]
- ▶ K3 surfaces [complete classification of D-equivalence by Mukai]
- ▶ Calabi-Yau threefolds [examples known by Borisov-Caldararu, Kuznetsov...]

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

- ▶ Mukai: if A abelian variety, then $\mathcal{D}^b(A) \simeq \mathcal{D}^b(\widehat{A})$
- ▶ Abelian varieties [complete classification of D-equivalence by Orlov]
- ▶ K3 surfaces [complete classification of D-equivalence by Mukai]
- ▶ Calabi-Yau threefolds [examples known by Borisov-Caldararu, Kuznetsov...]
- ▶ Kuznetsov: Homological Projective Duality

D-equivalence

All varieties smooth projective over a field k of characteristic zero.

Definition

- ▶ $\mathcal{D}^b(X) = \mathcal{D}^b(\text{Coh}(X))$, bounded derived category of coherent sheaves
- ▶ X and Y are called derived equivalent or D-equivalent if $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

Examples of $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y)$

- ▶ Mukai: if A abelian variety, then $\mathcal{D}^b(A) \simeq \mathcal{D}^b(\widehat{A})$
- ▶ Abelian varieties [complete classification of D-equivalence by Orlov]
- ▶ K3 surfaces [complete classification of D-equivalence by Mukai]
- ▶ Calabi-Yau threefolds [examples known by Borisov-Caldararu, Kuznetsov...]
- ▶ Kuznetsov: Homological Projective Duality
- ▶ Some flops [Bondal-Orlov, Bridgeland...]

D-equivalence and K-equivalence

D-equivalence and K-equivalence

D-equivalence and geometry

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence
- ▶ Big question: are Hodge numbers preserved under D-equivalence?

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence
- ▶ Big question: are Hodge numbers preserved under D-equivalence?

Definition

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence
- ▶ Big question: are Hodge numbers preserved under D-equivalence?

Definition

- ▶ X and Y are called K -equivalent if there exists a Z together with birational morphisms $f: Z \rightarrow X$, $g: Z \rightarrow Y$ such that $f^*(K_X) = g^*(K_Y)$.

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence
- ▶ Big question: are Hodge numbers preserved under D-equivalence?

Definition

- ▶ X and Y are called K -equivalent if there exists a Z together with birational morphisms $f: Z \rightarrow X$, $g: Z \rightarrow Y$ such that $f^*(K_X) = g^*(K_Y)$.

Conjecture (Kawamata)

If X and Y are birational then X and Y are D-equivalent if and only if they are K-equivalent.

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence
- ▶ Big question: are Hodge numbers preserved under D-equivalence?

Definition

- ▶ X and Y are called K -equivalent if there exists a Z together with birational morphisms $f: Z \rightarrow X$, $g: Z \rightarrow Y$ such that $f^*(K_X) = g^*(K_Y)$.

Conjecture (Kawamata)

If X and Y are birational then X and Y are D-equivalent if and only if they are K-equivalent.

- ▶ K-equivalence \implies D-equivalence is known in large generality

D-equivalence and K-equivalence

D-equivalence and geometry

- ▶ D-equivalence does not imply birational equivalence
- ▶ Big question: are Hodge numbers preserved under D-equivalence?

Definition

- ▶ X and Y are called K -equivalent if there exists a Z together with birational morphisms $f: Z \rightarrow X$, $g: Z \rightarrow Y$ such that $f^*(K_X) = g^*(K_Y)$.

Conjecture (Kawamata)

If X and Y are birational then X and Y are D-equivalent if and only if they are K-equivalent.

- ▶ K-equivalence \implies D-equivalence is known in large generality
- ▶ For instance, it is known in dimension 3 [Kawamata]

D-equivalence and moduli spaces

D-equivalence and moduli spaces

Moduli spaces

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

Theorem (Mukai)

Let X and Y be K3 surfaces. Then X and Y are D-equivalent if and only if Y is a 2-dimensional fine moduli space of sheaves on X :

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

Theorem (Mukai)

Let X and Y be K3 surfaces. Then X and Y are D-equivalent if and only if Y is a 2-dimensional fine moduli space of sheaves on X : $Y = M_X(r, H, s)$, $H^2 = 2rs$.

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

Theorem (Mukai)

Let X and Y be K3 surfaces. Then X and Y are D-equivalent if and only if Y is a 2-dimensional fine moduli space of sheaves on X : $Y = M_X(r, H, s)$, $H^2 = 2rs$.

Non-fine moduli spaces and the Brauer class

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

Theorem (Mukai)

Let X and Y be K3 surfaces. Then X and Y are D-equivalent if and only if Y is a 2-dimensional fine moduli space of sheaves on X : $Y = M_X(r, H, s)$, $H^2 = 2rs$.

Non-fine moduli spaces and the Brauer class

- ▶ Let Y be non-fine moduli space of sheaves on X (X, Y K3 surfaces)

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

Theorem (Mukai)

Let X and Y be K3 surfaces. Then X and Y are D-equivalent if and only if Y is a 2-dimensional fine moduli space of sheaves on X : $Y = M_X(r, H, s)$, $H^2 = 2rs$.

Non-fine moduli spaces and the Brauer class

- ▶ Let Y be non-fine moduli space of sheaves on X (X, Y K3 surfaces)
- ▶ Then the universal sheaf \mathcal{E} on $X \times Y$ is an α -twisted sheaf, $\alpha \in \text{Br}(Y)$

D-equivalence and moduli spaces

Moduli spaces

- ▶ Let Y be a fine moduli space of sheaves on X
- ▶ Let \mathcal{E} be the universal sheaf on $X \times Y$
- ▶ Then \mathcal{E} determines a functor $\mathcal{D}^b(Y) \rightarrow \mathcal{D}^b(X)$ (Fourier-Mukai transform)

Theorem (Mukai)

Let X and Y be K3 surfaces. Then X and Y are D-equivalent if and only if Y is a 2-dimensional fine moduli space of sheaves on X : $Y = M_X(r, H, s)$, $H^2 = 2rs$.

Non-fine moduli spaces and the Brauer class

- ▶ Let Y be non-fine moduli space of sheaves on X (X, Y K3 surfaces)
- ▶ Then the universal sheaf \mathcal{E} on $X \times Y$ is an α -twisted sheaf, $\alpha \in \text{Br}(Y)$
- ▶ $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y, \alpha)$ [Mukai, Caldararu...].

Introduction: Geometric meaning of D-equivalence

Grothendieck ring of varieties and L-equivalence

Quadrics, quadric fibrations and K3 surfaces

The Grothendieck ring $K_0(\text{Var}/k)$

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety
- ▶ Relations: $[X] = [Z] + [U]$ for any closed $Z \subset X$ with open complement U

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety
- ▶ Relations: $[X] = [Z] + [U]$ for any closed $Z \subset X$ with open complement U
- ▶ Product: $[X] \cdot [Y] = [X \times Y]$

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety
 - ▶ Relations: $[X] = [Z] + [U]$ for any closed $Z \subset X$ with open complement U
 - ▶ Product: $[X] \cdot [Y] = [X \times Y]$
-
- ▶ $\mathbb{L} := [\mathbb{A}^1]$ - the Lefschetz class (sometimes called the Tate class)

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety
- ▶ Relations: $[X] = [Z] + [U]$ for any closed $Z \subset X$ with open complement U
- ▶ Product: $[X] \cdot [Y] = [X \times Y]$

- ▶ $\mathbb{L} := [\mathbb{A}^1]$ - the Lefschetz class (sometimes called the Tate class)
- ▶ $[\mathbb{P}^n] = \sum_{k=0}^n [\mathbb{A}^k] = \sum_{k=0}^n \mathbb{L}^k$

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety
- ▶ Relations: $[X] = [Z] + [U]$ for any closed $Z \subset X$ with open complement U
- ▶ Product: $[X] \cdot [Y] = [X \times Y]$

▶ $\mathbb{L} := [\mathbb{A}^1]$ - the Lefschetz class (sometimes called the Tate class)

▶ $[\mathbb{P}^n] = \sum_{k=0}^n [\mathbb{A}^k] = \sum_{k=0}^n \mathbb{L}^k$

Example

The Grothendieck ring $K_0(\text{Var}/k)$

Definition

$K_0(\text{Var}/k)$ is the following ring:

- ▶ Generators: $[X]$, X/k quasi-projective variety
- ▶ Relations: $[X] = [Z] + [U]$ for any closed $Z \subset X$ with open complement U
- ▶ Product: $[X] \cdot [Y] = [X \times Y]$

- ▶ $\mathbb{L} := [\mathbb{A}^1]$ - the Lefschetz class (sometimes called the Tate class)
- ▶ $[\mathbb{P}^n] = \sum_{k=0}^n [\mathbb{A}^k] = \sum_{k=0}^n \mathbb{L}^k$

Example

If X and Y are related by a flop then $[X] = [Y]$.

Some properties of the Grothendieck ring

Simple properties

Some properties of the Grothendieck ring

Simple properties

- ▶ If $X \rightarrow B$ a Zarisky locally trivial fibration with fiber F , then

$$[X] = [F] \cdot [B]$$

Some properties of the Grothendieck ring

Simple properties

- ▶ If $X \rightarrow B$ a Zarisky locally trivial fibration with fiber F , then

$$[X] = [F] \cdot [B]$$

- ▶ If $Y = Bl_Z(X)$ is a blow up of a smooth $Z \subset X$, then

$$[Y] - [\mathbb{P}(N_{Z/X})] = [X] - [Z]$$

Some properties of the Grothendieck ring

Simple properties

- ▶ If $X \rightarrow B$ a Zarisky locally trivial fibration with fiber F , then

$$[X] = [F] \cdot [B]$$

- ▶ If $Y = Bl_Z(X)$ is a blow up of a smooth $Z \subset X$, then

$$[Y] - [\mathbb{P}(N_{Z/X})] = [X] - [Z]$$

Old open question: what is the geometric meaning of $[X] = [Y]$?

Some properties of the Grothendieck ring

Simple properties

- ▶ If $X \rightarrow B$ a Zarisky locally trivial fibration with fiber F , then

$$[X] = [F] \cdot [B]$$

- ▶ If $Y = Bl_Z(X)$ is a blow up of a smooth $Z \subset X$, then

$$[Y] - [\mathbb{P}(N_{Z/X})] = [X] - [Z]$$

Old open question: what is the geometric meaning of $[X] = [Y]$?

Theorem (Larsen-Lunts)

We have $[X] \equiv [Y] \pmod{\mathbb{L}}$ if and only if X and Y are stably birational.

Some properties of the Grothendieck ring

Simple properties

- ▶ If $X \rightarrow B$ a Zarisky locally trivial fibration with fiber F , then

$$[X] = [F] \cdot [B]$$

- ▶ If $Y = Bl_Z(X)$ is a blow up of a smooth $Z \subset X$, then

$$[Y] - [\mathbb{P}(N_{Z/X})] = [X] - [Z]$$

Old open question: what is the geometric meaning of $[X] = [Y]$?

Theorem (Larsen-Lunts)

We have $[X] \equiv [Y] \pmod{\mathbb{L}}$ if and only if X and Y are stably birational.

Corollary

If X, Y are non-uniruled (e.g. K3s or Calabi-Yau) such that $[X] \equiv [Y] \pmod{\mathbb{L}}$ then X and Y birational.

Zero divisors in the Grothendieck ring

Zero divisors in the Grothendieck ring

It has been known for a long time that $K_0(\text{Var}/k)$ is not a domain [Shioda, Poonen...],

Zero divisors in the Grothendieck ring

It has been known for a long time that $K_0(\text{Var}/k)$ is not a domain [Shioda, Poonen...], e.g. elliptic curves $E' \not\cong E''$ which satisfy

$$E \times E' \simeq E \times E''$$

for some E gives

$$[E]([E'] - [E'']) = 0$$

but both terms are nonzero.

Zero divisors in the Grothendieck ring

It has been known for a long time that $K_0(\text{Var}/k)$ is not a domain [Shioda, Poonen...], e.g. elliptic curves $E' \not\cong E''$ which satisfy

$$E \times E' \simeq E \times E''$$

for some E gives

$$[E]([E'] - [E'']) = 0$$

but both terms are nonzero.

Old question: is $\mathbb{L} = [\mathbb{A}^1] \in K_0(\text{Var}/k)$ a zero-divisor?

Zero divisors in the Grothendieck ring

It has been known for a long time that $K_0(\text{Var}/k)$ is not a domain [Shioda, Poonen...], e.g. elliptic curves $E' \not\cong E''$ which satisfy

$$E \times E' \simeq E \times E''$$

for some E gives

$$[E]([E'] - [E'']) = 0$$

but both terms are nonzero.

Old question: is $\mathbb{L} = [\mathbb{A}^1] \in K_0(\text{Var}/k)$ a zero-divisor?

- ▶ Galkin-S. 2014: if \mathbb{L} is not a zero divisor, then a very general complex cubic fourfold X is irrational (and a sufficient condition for irrationality in terms of the Fano variety of lines $F(X)$)

Zero divisors in the Grothendieck ring

It has been known for a long time that $K_0(\text{Var}/k)$ is not a domain [Shioda, Poonen...], e.g. elliptic curves $E' \not\cong E''$ which satisfy

$$E \times E' \simeq E \times E''$$

for some E gives

$$[E]([E'] - [E'']) = 0$$

but both terms are nonzero.

Old question: is $\mathbb{L} = [\mathbb{A}^1] \in K_0(\text{Var}/k)$ a zero-divisor?

- ▶ Galkin-S. 2014: if \mathbb{L} is not a zero divisor, then a very general complex cubic fourfold X is irrational (and a sufficient condition for irrationality in terms of the Fano variety of lines $F(X)$)
- ▶ Borisov 2014 (improved by Martin 2016): $\mathbb{L}^6([X] - [Y]) = 0$ for Calabi-Yau threefolds X and Y in the Pfaffian-Grassmannian correspondence

L-equivalence

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

Known examples

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

Known examples

- ▶ Calabi-Yau threefolds: Borisov (Pfaffian-Grassmannian correspondence); Ito–Miura–Okawa–Ueda (G^2 geometry)

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

Known examples

- ▶ Calabi-Yau threefolds: Borisov (Pfaffian-Grassmannian correspondence); Ito–Miura–Okawa–Ueda (G^2 geometry)
- ▶ K3 surfaces: Kuznetsov-S. (degrees 8 and 2); Ito–Miura–Okawa–Ueda, Hassett-Lai (degrees 12 and 12); Kuznetsov-S. (degree 16 and 4: work in progress)

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

Known examples

- ▶ Calabi-Yau threefolds: Borisov (Pfaffian-Grassmannian correspondence); Ito–Miura–Okawa–Ueda (G^2 geometry)
- ▶ K3 surfaces: Kuznetsov-S. (degrees 8 and 2); Ito–Miura–Okawa–Ueda, Hassett-Lai (degrees 12 and 12); Kuznetsov-S. (degree 16 and 4: work in progress)

In all these examples pairs of varieties X and Y are D-equivalent.

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

Known examples

- ▶ Calabi-Yau threefolds: Borisov (Pfaffian-Grassmannian correspondence); Ito–Miura–Okawa–Ueda (G^2 geometry)
- ▶ K3 surfaces: Kuznetsov-S. (degrees 8 and 2); Ito–Miura–Okawa–Ueda, Hassett-Lai (degrees 12 and 12); Kuznetsov-S. (degree 16 and 4: work in progress)

In all these examples pairs of varieties X and Y are D-equivalent.

Conjecture

D-equivalence of X and Y implies their L-equivalence.

L-equivalence

Definition

We call X and Y L-equivalent if $\mathbb{L}^n([X] - [Y]) = 0$ for some $n \geq 1$.

Known examples

- ▶ Calabi-Yau threefolds: Borisov (Pfaffian-Grassmannian correspondence); Ito–Miura–Okawa–Ueda (G^2 geometry)
- ▶ K3 surfaces: Kuznetsov-S. (degrees 8 and 2); Ito–Miura–Okawa–Ueda, Hassett-Lai (degrees 12 and 12); Kuznetsov-S. (degree 16 and 4: work in progress)

In all these examples pairs of varieties X and Y are D-equivalent.

Conjecture

D-equivalence of X and Y implies their L-equivalence.

It is easy to see that L-equivalent varieties have the same Hodge numbers, so this would imply invariance of Hodge numbers under D-equivalence.

Introduction: Geometric meaning of D-equivalence

Grothendieck ring of varieties and L-equivalence

Quadrics, quadric fibrations and K3 surfaces

Warm up: elliptic curves

Warm up: elliptic curves

Construction

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$
- ▶ $f(\lambda) = \det(Q_\lambda)$ has degree 4

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$
- ▶ $f(\lambda) = \det(Q_\lambda)$ has degree 4
- ▶ $Z \subset \mathbb{P}^1$ four points corresponding to singular quadrics

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$
- ▶ $f(\lambda) = \det(Q_\lambda)$ has degree 4
- ▶ $Z \subset \mathbb{P}^1$ four points corresponding to singular quadrics
- ▶ $Y \rightarrow \mathbb{P}^1$ double cover ramified in Z ; Y is an elliptic curve.

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$
- ▶ $f(\lambda) = \det(Q_\lambda)$ has degree 4
- ▶ $Z \subset \mathbb{P}^1$ four points corresponding to singular quadrics
- ▶ $Y \rightarrow \mathbb{P}^1$ double cover ramified in Z ; Y is an elliptic curve.

Properties of X and Y

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$
- ▶ $f(\lambda) = \det(Q_\lambda)$ has degree 4
- ▶ $Z \subset \mathbb{P}^1$ four points corresponding to singular quadrics
- ▶ $Y \rightarrow \mathbb{P}^1$ double cover ramified in Z ; Y is an elliptic curve.

Properties of X and Y

- ▶ Over an algebraically closed field, X and Y are D-equivalent, L-equivalent, and in fact isomorphic.

Warm up: elliptic curves

Construction

- ▶ $X = Q_1 \cap Q_2 \subset \mathbb{P}^3$ smooth complete intersection of two quadrics; X is an elliptic curve.
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2$ pencil of quadrics: $Q \rightarrow \mathbb{P}^1$
- ▶ $f(\lambda) = \det(Q_\lambda)$ has degree 4
- ▶ $Z \subset \mathbb{P}^1$ four points corresponding to singular quadrics
- ▶ $Y \rightarrow \mathbb{P}^1$ double cover ramified in Z ; Y is an elliptic curve.

Properties of X and Y

- ▶ Over an algebraically closed field, X and Y are D-equivalent, L-equivalent, and in fact isomorphic.
- ▶ Over non-algebraically closed field, X and Y don't have to be isomorphic, but are unlikely to be D-equivalent or L-equivalent either

Homological projective duality for K3 surfaces of degree 8

Homological projective duality for K3 surfaces of degree 8

Setup

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$
- ▶ $C = \det(Q_\lambda) \subset \mathbb{P}^2$, a smooth sextic

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$
- ▶ $C = \det(Q_\lambda) \subset \mathbb{P}^2$, a smooth sextic
- ▶ $Y \rightarrow \mathbb{P}^2$ double covering ramified in C

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$
- ▶ $C = \det(Q_\lambda) \subset \mathbb{P}^2$, a smooth sextic
- ▶ $Y \rightarrow \mathbb{P}^2$ double covering ramified in C
- ▶ $\alpha_Y \in \text{Br}(Y)$ determined by the Clifford algebra of Q_λ

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$
- ▶ $C = \det(Q_\lambda) \subset \mathbb{P}^2$, a smooth sextic
- ▶ $Y \rightarrow \mathbb{P}^2$ double covering ramified in C
- ▶ $\alpha_Y \in \text{Br}(Y)$ determined by the Clifford algebra of Q_λ

We have constructed: a K3 surface X of degree 8 and a K3 surface Y of degree 2 that are in a certain sense dual to each other.

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$
- ▶ $C = \det(Q_\lambda) \subset \mathbb{P}^2$, a smooth sextic
- ▶ $Y \rightarrow \mathbb{P}^2$ double covering ramified in C
- ▶ $\alpha_Y \in \text{Br}(Y)$ determined by the Clifford algebra of Q_λ

We have constructed: a K3 surface X of degree 8 and a K3 surface Y of degree 2 that are in a certain sense dual to each other.

Theorem (Mukai, Caldararu, Kuznetsov)

There is an equivalence $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y, \alpha_Y)$. In particular, if $\alpha_Y = 0$, then X and Y are D-equivalent.

Homological projective duality for K3 surfaces of degree 8

Setup

- ▶ $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$, smooth complete intersection of 3 quadrics
- ▶ $Q_\lambda = \lambda_1 Q_1 + \lambda_2 Q_2 + \lambda_3 Q_3$ net of quadrics ($\lambda \in \mathbb{P}^2$): $Q \rightarrow \mathbb{P}^2$
- ▶ $C = \det(Q_\lambda) \subset \mathbb{P}^2$, a smooth sextic
- ▶ $Y \rightarrow \mathbb{P}^2$ double covering ramified in C
- ▶ $\alpha_Y \in \text{Br}(Y)$ determined by the Clifford algebra of Q_λ

We have constructed: a K3 surface X of degree 8 and a K3 surface Y of degree 2 that are in a certain sense dual to each other.

Theorem (Mukai, Caldararu, Kuznetsov)

There is an equivalence $\mathcal{D}^b(X) \simeq \mathcal{D}^b(Y, \alpha_Y)$. In particular, if $\alpha_Y = 0$, then X and Y are D-equivalent.

Mukai's approach: $Y = M_X(2, H, 2)$, the non-fine moduli space of spinor bundles on X and $\alpha_Y = 0$ if and only if the moduli space is fine.

Main Theorem

Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$.

Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

In the rest of the talk we are proving this Theorem.

Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

In the rest of the talk we are proving this Theorem.

Idea: consider the universal quadric $Q \subset \mathbb{P}^5 \times \mathbb{P}^2$ and compute its class $[Q]$ in two ways to get a relation between $[X]$ and $[Y]$.


Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

In the rest of the talk we are proving this Theorem.

Idea: consider the universal quadric $Q \subset \mathbb{P}^5 \times \mathbb{P}^2$ and compute its class $[Q]$ in two ways to get a relation between $[X]$ and $[Y]$.




Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

In the rest of the talk we are proving this Theorem.

Idea: consider the universal quadric $Q \subset \mathbb{P}^5 \times \mathbb{P}^2$ and compute its class $[Q]$ in two ways to get a relation between $[X]$ and $[Y]$.



- ▶ p is piece-wise locally trivial so can relate $[H]$ to $[\mathbb{P}^5]$ and $[X]$


Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

In the rest of the talk we are proving this Theorem.

Idea: consider the universal quadric $Q \subset \mathbb{P}^5 \times \mathbb{P}^2$ and compute its class $[Q]$ in two ways to get a relation between $[X]$ and $[Y]$.



- ▶ p is piece-wise locally trivial so can relate $[H]$ to $[\mathbb{P}^5]$ and $[X]$
- ▶ Fibers of q over $\lambda \in \mathbb{P}^2$ are quadrics Q_λ ; q is NOT locally trivial


Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

In the rest of the talk we are proving this Theorem.

Idea: consider the universal quadric $Q \subset \mathbb{P}^5 \times \mathbb{P}^2$ and compute its class $[Q]$ in two ways to get a relation between $[X]$ and $[Y]$.



- ▶ p is piece-wise locally trivial so can relate $[H]$ to $[\mathbb{P}^5]$ and $[X]$
- ▶ Fibers of q over $\lambda \in \mathbb{P}^2$ are quadrics Q_λ ; q is NOT locally trivial
- ▶ Use hyperbolic reduction of quadrics to relate $[H]$ to the double cover Y

Hyperbolic reduction

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Geometric meaning

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Geometric meaning

- ▶ Let $Q \subset \mathbb{P}(V)$ be the quadric corresponding to (V, q)

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Geometric meaning

- ▶ Let $Q \subset \mathbb{P}(V)$ be the quadric corresponding to (V, q)
- ▶ Let $x \in Q$ be the point corresponding to $L = \langle v \rangle \subset V$

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Geometric meaning

- ▶ Let $Q \subset \mathbb{P}(V)$ be the quadric corresponding to (V, q)
- ▶ Let $x \in Q$ be the point corresponding to $L = \langle v \rangle \subset V$
- ▶ Projecting from $x \in Q$ is a birational map from Q to $\mathbb{P}(V/L)$

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Geometric meaning

- ▶ Let $Q \subset \mathbb{P}(V)$ be the quadric corresponding to (V, q)
- ▶ Let $x \in Q$ be the point corresponding to $L = \langle v \rangle \subset V$
- ▶ Projecting from $x \in Q$ is a birational map from Q to $\mathbb{P}(V/L)$
- ▶ Projection contracts the set of lines on Q through x to a quadric $\bar{Q} \subset \mathbb{P}(V/L)$ corresponding to (\bar{V}, \bar{q})

Hyperbolic reduction

Hyperbolic reduction for a quadratic form

- ▶ Let (V, q) be a nondegenerate quadratic form
- ▶ $v \in V$ isotropic vector (that is $q(v) = 0$)
- ▶ Let $\bar{V} = \langle v \rangle^\perp / \langle v \rangle$ and let \bar{q} be the induced quadratic form on \bar{V}
- ▶ Then $\dim(\bar{V}) = \dim(V) - 2$ and $V = \bar{V} \perp \langle 1, -1 \rangle$.
- ▶ We call (\bar{V}, \bar{q}) the hyperbolic reduction of (V, q)

Geometric meaning

- ▶ Let $Q \subset \mathbb{P}(V)$ be the quadric corresponding to (V, q)
- ▶ Let $x \in Q$ be the point corresponding to $L = \langle v \rangle \subset V$
- ▶ Projecting from $x \in Q$ is a birational map from Q to $\mathbb{P}(V/L)$
- ▶ Projection contracts the set of lines on Q through x to a quadric $\bar{Q} \subset \mathbb{P}(V/L)$ corresponding to (\bar{V}, \bar{q})
- ▶ We get a relation $[Q] = 1 + \mathbb{L}^{\dim(Q)} + \mathbb{L}[\bar{Q}] \in K_0(\text{Var}/k)$.

Quadric fibrations with a smooth section

Quadric fibrations with a smooth section

Lemma

If $Q \rightarrow S$ is a quadric fibration of relative dimension n and $s: S \rightarrow Q$ is a smooth section, then

$$[Q] = (1 + \mathbb{L}^n)[S] + \mathbb{L}[\overline{Q}]$$

where \overline{Q} is the fiberwise hyperbolic reduction of Q .

Quadric fibrations with a smooth section

Lemma

If $Q \rightarrow S$ is a quadric fibration of relative dimension n and $s: S \rightarrow Q$ is a smooth section, then

$$[Q] = (1 + \mathbb{L}^n)[S] + \mathbb{L}[\overline{Q}]$$

where \overline{Q} is the fiberwise hyperbolic reduction of Q .

Proof.

Project the quadric fibration from the section s and resolve the indeterminacy locus of the projection. □

Quadric fibrations with a smooth section

Lemma

If $Q \rightarrow S$ is a quadric fibration of relative dimension n and $s: S \rightarrow Q$ is a smooth section, then

$$[Q] = (1 + \mathbb{L}^n)[S] + \mathbb{L}[\overline{Q}]$$

where \overline{Q} is the fiberwise hyperbolic reduction of Q .

Proof.

Project the quadric fibration from the section s and resolve the indeterminacy locus of the projection. □

Example

Let $Q \rightarrow \mathbb{P}^m$ be a linear system of quadrics in \mathbb{P}^{n+1} . Let X be the base locus of this system. Any smooth point $x \in X$ determines a section of Q and the Lemma allows us to relate $[Q]$ and $[\overline{Q}]$.

Quadric fibrations with a smooth section

Lemma

If $Q \rightarrow S$ is a quadric fibration of relative dimension n and $s: S \rightarrow Q$ is a smooth section, then

$$[Q] = (1 + \mathbb{L}^n)[S] + \mathbb{L}[\overline{Q}]$$

where \overline{Q} is the fiberwise hyperbolic reduction of Q .

Proof.

Project the quadric fibration from the section s and resolve the indeterminacy locus of the projection. □

Example

Let $Q \rightarrow \mathbb{P}^m$ be a linear system of quadrics in \mathbb{P}^{n+1} . Let X be the base locus of this system. Any smooth point $x \in X$ determines a section of Q and the Lemma allows us to relate $[Q]$ and $[\overline{Q}]$.

This applies in particular to our K3 surface $X = Q_1 \cap Q_2 \cap Q_3$ and gives us a quadric fibration $\overline{Q} \rightarrow \mathbb{P}^2$ of relative dimension 2 with the same Brauer class α_Y .

Quadric fibrations of relative dimension two

Quadric fibrations of relative dimension two

Setup

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations
- ▶ $Y \rightarrow S$ discriminant double cover of $Q \rightarrow S$, $\alpha_Y \in \text{Br}(Y)$

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations
- ▶ $Y \rightarrow S$ discriminant double cover of $Q \rightarrow S$, $\alpha_Y \in \text{Br}(Y)$
- ▶ Hyperbolic reduction \overline{Q} (if sections exist) is canonically identified with Y

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations
- ▶ $Y \rightarrow S$ discriminant double cover of $Q \rightarrow S$, $\alpha_Y \in \text{Br}(Y)$
- ▶ Hyperbolic reduction \overline{Q} (if sections exist) is canonically identified with Y

Lemma

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations
- ▶ $Y \rightarrow S$ discriminant double cover of $Q \rightarrow S$, $\alpha_Y \in \text{Br}(Y)$
- ▶ Hyperbolic reduction \overline{Q} (if sections exist) is canonically identified with Y

Lemma

- ▶ $\alpha_Y = 0 \iff Q \rightarrow S$ has a rational multisection of odd degree.

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations
- ▶ $Y \rightarrow S$ discriminant double cover of $Q \rightarrow S$, $\alpha_Y \in \text{Br}(Y)$
- ▶ Hyperbolic reduction \overline{Q} (if sections exist) is canonically identified with Y

Lemma

- ▶ $\alpha_Y = 0 \iff Q \rightarrow S$ has a rational multisection of odd degree.
- ▶ If $\alpha_Y = 0$, then $[Q] = (1 + \mathbb{L}^2)[S] + \mathbb{L}[Y]$.

Quadric fibrations of relative dimension two

Setup

- ▶ $Q \rightarrow S$ a quadric fibration of relative dimension 2 with nodal degenerations
- ▶ $Y \rightarrow S$ discriminant double cover of $Q \rightarrow S$, $\alpha_Y \in \text{Br}(Y)$
- ▶ Hyperbolic reduction \overline{Q} (if sections exist) is canonically identified with Y

Lemma

- ▶ $\alpha_Y = 0 \iff Q \rightarrow S$ has a rational multisection of odd degree.
- ▶ If $\alpha_Y = 0$, then $[Q] = (1 + \mathbb{L}^2)[S] + \mathbb{L}[Y]$.

Example

In the case of K3 surface $X = Q_1 \cap Q_2 \cap Q_3 \subset \mathbb{P}^5$ vanishing $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree, and Y is the dual K3 surface of degree 2.

Proof of the Main Theorem

Proof of the Main Theorem

Theorem (Kuznetsov-S.)


If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Proof of the Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Proof




Proof of the Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Proof




- p is piece-wise locally trivial: $[Q] = [\mathbb{P}^5][\mathbb{P}^1] + \mathbb{L}^2[X]$

Proof of the Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Proof



- ▶ p is piece-wise locally trivial: $[Q] = [\mathbb{P}^5][\mathbb{P}^1] + \mathbb{L}^2[X]$
- ▶ First hyperbolic reduction for q and a choice of a point $x \in X$:


$$[Q] = [\mathbb{P}^2](1 + \mathbb{L}^4) + \mathbb{L}[\overline{Q}]$$

Proof of the Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Proof



- ▶ p is piece-wise locally trivial: $[Q] = [\mathbb{P}^5][\mathbb{P}^1] + \mathbb{L}^2[X]$
- ▶ First hyperbolic reduction for q and a choice of a point $x \in X$:

$$[Q] = [\mathbb{P}^2](1 + \mathbb{L}^4) + \mathbb{L}[\overline{Q}]$$


- ▶ Second hyperbolic reduction ($\alpha_Y = 0$): $[\overline{Q}] = [\mathbb{P}^2](1 + \mathbb{L}^2) + \mathbb{L}[Y]$

Proof of the Main Theorem

Theorem (Kuznetsov-S.)

If X and Y are dual K3 surfaces of degree 8 and 2 respectively such that $\alpha_Y = 0$, then $\mathbb{L}^2([X] - [Y]) = 0$. For general such X and Y we have $[X] \neq [Y]$.

Proof



- ▶ p is piece-wise locally trivial: $[Q] = [\mathbb{P}^5][\mathbb{P}^1] + \mathbb{L}^2[X]$
- ▶ First hyperbolic reduction for q and a choice of a point $x \in X$:

$$[Q] = [\mathbb{P}^2](1 + \mathbb{L}^4) + \mathbb{L}[\overline{Q}]$$

- ▶ Second hyperbolic reduction ($\alpha_Y = 0$): $[\overline{Q}] = [\mathbb{P}^2](1 + \mathbb{L}^2) + \mathbb{L}[Y]$
- ▶ Finally: canceling matching terms gives $\mathbb{L}^2[X] = \mathbb{L}^2[Y]$

Proof continued: showing that X and Y are not always isomorphic

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]
- ▶ There are infinitely many d such that $\alpha_Y = 0$ but $X \not\simeq Y$: for instance take $d = k^2$, an odd square; and this finishes the proof!

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]
- ▶ There are infinitely many d such that $\alpha_Y = 0$ but $X \not\simeq Y$: for instance take $d = k^2$, an odd square; and this finishes the proof!

Examples

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]
- ▶ There are infinitely many d such that $\alpha_Y = 0$ but $X \not\simeq Y$: for instance take $d = k^2$, an odd square; and this finishes the proof!

Examples

- ▶ X contains a line $\implies \alpha_Y = 0$, $X \simeq Y$ (classical geometric construction!)

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]
- ▶ There are infinitely many d such that $\alpha_Y = 0$ but $X \not\simeq Y$: for instance take $d = k^2$, an odd square; and this finishes the proof!

Examples

- ▶ X contains a line $\implies \alpha_Y = 0$, $X \simeq Y$ (classical geometric construction!)
- ▶ X contains a conic $\implies \alpha_Y \neq 0$ (and $X \not\simeq Y$ generically)

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]
- ▶ There are infinitely many d such that $\alpha_Y = 0$ but $X \not\simeq Y$: for instance take $d = k^2$, an odd square; and this finishes the proof!

Examples

- ▶ X contains a line $\implies \alpha_Y = 0$, $X \simeq Y$ (classical geometric construction!)
- ▶ X contains a conic $\implies \alpha_Y \neq 0$ (and $X \not\simeq Y$ generically)
- ▶ X contains a twisted cubic $\implies \alpha_Y = 0$, $X \not\simeq Y$ generically

Proof continued: showing that X and Y are not always isomorphic

- ▶ Recall: $\alpha_Y = 0$ is equivalent to existence of a curve $C \subset X$ of odd degree
- ▶ This condition singles out a countable union of divisors $\mathcal{D}_d \subset \mathcal{M}_8$ in the moduli space of K3 surfaces X of degree 8, namely: $d \equiv 1 \pmod{8}$
- ▶ Here \mathcal{D}_d parametrizes K3 surfaces of degree 8 and Picard rank 2 with discriminant $-d$
- ▶ The condition $X \simeq Y$ singles out the divisors \mathcal{D}_d such that $a^2 - db^2 = \pm 8$ has a solution [Madonna-Nikulin]
- ▶ There are infinitely many d such that $\alpha_Y = 0$ but $X \not\simeq Y$: for instance take $d = k^2$, an odd square; and this finishes the proof!

Examples

- ▶ X contains a line $\implies \alpha_Y = 0$, $X \simeq Y$ (classical geometric construction!)
- ▶ X contains a conic $\implies \alpha_Y \neq 0$ (and $X \not\simeq Y$ generically)
- ▶ X contains a twisted cubic $\implies \alpha_Y = 0$, $X \not\simeq Y$ generically

Remark: refining the argument one can show that $\alpha_Y = 0 \implies \mathbb{L}([X] - [Y]) = 0$.

Homework

Four open questions which may be accessible

Homework

Four open questions which may be accessible

1. Can curves be L-equivalent?

Homework

Four open questions which may be accessible

1. Can curves be L-equivalent?
2. Can abelian varieties be L-equivalent, e.g. what about A and \widehat{A} ?

Homework


Four open questions which may be accessible

1. Can curves be L-equivalent?
2. Can abelian varieties be L-equivalent, e.g. what about A and \widehat{A} ?
3. Prove that D-equivalence implies L-equivalence for K3 surfaces in general

Homework

Four open questions which may be accessible

1. Can curves be L-equivalent?
2. Can abelian varieties be L-equivalent, e.g. what about A and \widehat{A} ?
3. Prove that D-equivalence implies L-equivalence for K3 surfaces in general
4. How to describe the kernel $\text{Ker}(K_0(\text{Var}/k) \rightarrow K_0(\text{Var}/k)[\mathbb{L}^{-1}])$? Is it generated by $[X] - [Y]$ where X and Y are L-equivalent?



THE END